arXiv:hep-th/9405104v3 24 May 1994

SPhT/94-060

A lecture on the Calogero-Sutherland models.

V. Pasquier

*Service de Physique Théorique de Saclay *,*
F-91191 Gif sur Yvette Cedex, France

In these lectures, I review some recent results on the Calogero-Sutherland model and the Haldane Shatry-chain. The list of topics I cover are the following: 1) The Calogero-Sutherland Hamiltonien and fractional stastistics. The form factor of the density operator. 2) The Dunkl operators and their relations with monodromy matrices, Yangians and affine-Hecke algebras. 3) The Haldane-Shastry chain in connection with the Calogero-Sutherland Hamiltonian at a specific coupling constant.

05/94

* Laboratoire de la Direction des Sciences de la Matière du Commissariat à l'Energie Atomique.

1. INTRODUCTION

The Calogero-Sutherland model has recently attracted some attention mainly because, in spite of its simplicity, it yields nontrivial results which contribute to shape our understanding of fractional statistics [1]. Its most remarkable property is that its ground state is given by a Jastrow wave-function which is the one dimensional analogue of the Laughlin wave-function. The excitations are described by quasiparticles which carry a fraction of the quantum numbers of the fundamental particles. The wave functions are simple enough to allow the computation of physical quantities such as the dynamical correlation functions. Recently, it has become clear that the spin version of this model is closely related to the known spin chains such as the XXX chain. The wave functions of the spin models can be obtained as simply as those of the Calogero-Sutherland model by diagonalising simple differential operators known as the Dunkl operators [2].

In these lectures, I restrict to the simplest models and I introduce some techniques useful in their study. These lectures are based on work done in collaboration with D.Bernard, M.Gaudin and D.Haldane [3] and with F.Lesage and D.Serban [4].

In the first part, I review Sutherlands method to diagonalise the Calogero-Sutherland Hamiltonian [5]. I then give the expression of the form factor of the density operator. On this example, I describe the fractional character of the quasiparticles which propagate in the intermediate states.

In the second part, I review the spin generalisation of the Calogero-Sutherland model and I diagonalise their Hamiltonian using the Dunkl operators. I then exhibit a representation of the Yangian which commutes with the Hamiltonian. This is achieved by "quantizing" the spectral parameters of a monodromy matrix obeying the Yang-Baxter equation. The consistancy of this procedure requires that the spectral parameters (the Dunkl operators) obey the defining relations of a affine-Hecke-algebra. Finally, I obtain a representation of this algebra which degenerates to the Dunkl operators using some operators defined by Yang in his study of the δ-interacting gas [6].

In the last part, I review the long range interacting spin model known as the Haldane-Shastry chain [7][8]. Although closely related to the Calogero-Sutherland model, this chain is more difficult to study because the Dunkl operators cannot be used to obtain the wave-functions in a straightforward way. I exhibit a representation of the affine Hecke algebra in terms of parameters. This representation becomes reducible at the special point where it commutes with the Haldane-Shastry Hamiltonian. I then use a correspondance with the

Calogero-Sutherland models to obtain the eigenvalues and some of the eigenvectors of this Hamiltonian.

The most unphysical feature of the models discussed here is the long range character of the particle particle interaction which behaves as $1/x^2$ in the thermodynamical limit. More realistic models with short range potential ($1/\sinh(x)^2$) can be defined [9] but are more difficult to study (see [10] for some recent progress). Some aspects of the conformal limit of the Haldane-Shastry chain are also discussed in [11].

2. THE CALOGERO-SUTHERLAND HAMILTONIAN

This model describes particles on a circle interacting with a long range potential [3,4]. The positions of the particles are denoted by x_i, $1 \leq i \leq N$, $0 \leq x_i \leq L$ and the total momentum and Hamiltonian which give their dynamics are respectively given by:

$$\begin{aligned} P &= \sum_{j=1}^{N} \frac{1}{i} \frac{\mathrm{d}}{\mathrm{d}x_j} \\ H &= -\sum_{j=1}^{N} \frac{1}{2} \frac{\mathrm{d}^2}{\mathrm{d}x_j^2} + \beta(\beta-1)\frac{\pi^2}{L^2} \sum_{i<j} \frac{1}{\sin^2\left(\frac{\pi}{L}\left(x_i - x_j\right)\right)} \end{aligned} \tag{2.1}$$

From now on, we shall work on the unit circle and set $\theta_j = 2\pi \frac{x_j}{L}$ and $z_j = \mathrm{e}^{i\theta_j}$. The wave functions solution of the equation $H\psi = E\psi$ can be given the following structure:

$$\psi(\theta) = \phi(\theta)\Delta^{\beta}(\theta) \tag{2.2}$$

with

$$\Delta(\theta) = \prod_{i<j} \sin\left(\frac{\theta_i - \theta_j}{2}\right) \tag{2.3}$$

Φ is a symmetric polynomial in the variables $z_j = \mathrm{e}^{i\theta_j}$ and z_j^{-1}.

To understand the simplicity of the spectrum, one must write the effective Hamiltonian $\tilde{H} = \Delta^{-\beta} H \Delta^{\beta}$ acting on ϕ.
Following Sutherland [5], we obtain:

$$\tilde{H} = \sum_{j=1}^{N} \left(z_j \frac{\partial}{\partial z_j}\right)^2 + \beta \sum_{i \neq j} \frac{z_i + z_j}{z_i - z_j} \left(z_i \frac{\partial}{\partial z_i} - z_j \frac{\partial}{\partial z_j}\right) \tag{2.4}$$

The remarkable property of this Hamiltonian is that acting upon symmetric polynomials of a given homogeneity in the variables z_j it is realised as a triangular matrix. So, it's

eigenvalues can be read on the diagonal of the matrix and there are simple algorithms to find the eigenvectors.

More precisely, let us define the following basis of symmetric polynomials indexed by a partition: $\lambda_1 \geq \lambda_2 ... \geq \lambda_N \geq 0$.

$$m_{\{\lambda\}} = \sum z^{\lambda} \tag{2.5}$$

which is the sum over distinct permutations of the monomial $z_1^{\lambda_1} z_2^{\lambda_2} ... z_N^{\lambda_N}$.

First, it is easy to see that the subspace of polynomials of a given homogeneity ($\mid \lambda \mid = \sum_{i=1}^{N} \lambda_i$) is preserved. Then, inside this subspace, we can define an order on the partition by saying that $\lambda \geq \mu$ if $\lambda_1 \geq \mu_1$, $\lambda_1 + \lambda_2 \geq \mu_1 + \mu_2, ..., \lambda_1 + ... + \lambda_N \geq \mu_1 + \mu_2 + ... + \mu_N$. It follows from Sutherland's argument that:

$$\tilde{H} m_{\lambda} = \sum_{\mu \leq \lambda} C_{\lambda\mu} m_{\mu} \tag{2.6}$$

that is to say $\tilde{H}$ is a triangular matrix. The eigenvalues are given by the diagonal elements:

$$E_{\lambda} = C_{\lambda\lambda} = \sum_{i=1}^{N} \left(\lambda_i + (N-i)\beta\right)^2 - \left((N-i)\beta\right)^2 \tag{2.7}$$

The momentum is given by:

$$P_{\lambda} = \sum_{i=1}^{N} \left(\lambda_i + (N-i)\beta\right) \tag{2.8}$$

This set of states is not complete, but the complete set can be obtained by multiplying these wave functions by $(z_1 z_2 ... z_N)^p$ where p is a positive or negative integer. Taking this into account, a complete basis is given in terms of N momenta $k_i = \lambda_i + (N-i)\beta + p$ and the energy and momentum are given by:

$$\begin{aligned} P \left| k_1 ... k_N \right\rangle &= \left(\sum_{i=1}^{N} k_i \right) \left| k_1 ... k_N \right\rangle \\ H \left| k_1 ... k_N \right\rangle &= \left(\sum_{i=1}^{N} k_i^2 \right) \left| k_1 ... k_N \right\rangle \end{aligned} \tag{2.9}$$

This looks like a free particle spectrum, in particular for $\beta = 0, 1$, the Hamiltonian is a pure kinetic energy term and it reduces to the usual boson and fermion description of the

states. To understand the difference, let us consider the case of integer values of β. One sees from the expression of the k_i that they are integers and obey the constraint:

$$k_i - k_{i+1} \geq \beta \tag{2.10}$$

If $\beta = 1$, this is just the Pauli-exclusion principle, but if $\beta = 2, 3, ...$ the Pauli principle is replaced by a stronger exclusion principle.

2.1. Particle hole excitations

We shall now describe the excitations of the ground state of such a system and show that they can naturally be described in terms of quasiparticles. To illustrate this, we shall consider the matrix elements (form factor):

$$\langle \alpha \mid \rho(x) \mid 0 \rangle \tag{2.11}$$

of the density operator

$$\rho(x) = \sum_{i=1}^{N} \delta(\theta_i - x) \tag{2.12}$$

where the state $\mid 0 >$ stands for the ground state and $\mid \alpha >$ for some excited state. The results which we present here were motivated by the evaluation of the density density correlation function obtained by Simons, Lee and Altshuler in the matrix models [12] (see also [1]).

The question we address is for which states $\mid \alpha >$ is the matrix element non zero?

In the case $\beta = 1$ which corresponds to free fermions, it is easy to answer using second-quantization arguments. The operator $\rho(x)$ can be represented as $\psi^+(x)\psi(x)$ where $\psi(x)$ is a fermion field. Acting on the vacuum, the operator $\psi(x)$ anihilates a particle from the Fermi sea ($\mid k \mid \leq k_F$ where k_F is the Fermi momentum) and the operator $\psi^+(x)$ creates a particle out of the Fermi sea. The states $\mid \alpha >$ are therefore described by $(N-1)$ momenta k_i less than the Fermi momentum k_F and 1 momentum k_N larger than k_F or equivalently in terms of a hole of momentum less than k_F and a particle of momentum larger than k_F.

For β arbitrary integer, the calculation is based on the properties of the wave functions of $\tilde{H}$ (the Jack polynomials) established by Macdonald and Stanley [13][14]. The final result is however simple enough to be described here. Let us view the ground state as a Fermi-sea filled with N particles subject to the constraint $|k_i - k_j| \geq \beta$:

$$k_i = \beta \left(i - \frac{N+1}{2} \right), \qquad 1 \leq i \leq N \tag{2.13}$$

The Fermi momentum is given by the largest value of k_i :

$$k_F = \beta \frac{N-1}{2}. \tag{2.14}$$

This is represented on Figure 1 a) for $\beta = 2$, $N = 5$. The ones stand for the integers k_i occupied by a particle and the 0 for the remaining integers.



Fig. 1: The Fermi sea

It turns out that the states $\mid \alpha >$ which propagate are those for which $(N-1)$ momenta are inside the Fermi-sea $|k_i| \leq k_F$ and one momentum is out $k_N > k_F$ (see Figure 1 b)). Such states can be completely described by β holes of momentum less than k_F and a particle of momentum larger than k_F. A hole corresponds to a sequence of β consecutive integers not occupied by a momentum (represented by a cross on the Figure).

The outcome of the computation [4] (These results have also been conjectured in [15] and [16]) is that these holes are eigenstates of a Calogero-Sutherland Hamiltonian with coupling constant β^{-1} and have a mass equal to $\frac{1}{\beta}$ the mass of the particles. In the thermodynamic limit the form factor is equal to :

$$\langle \alpha \mid \rho(0) \mid 0 \rangle = (w^2-1)^{\frac{\beta-1}{2}} \prod_{i=1}^{\beta} (1-v_i^2)^{\frac{\beta^{-1}-1}{2}} \prod_{i<j} |v_i - v_j|^{1/\beta} \frac{(\Sigma_i v_i - \beta w)}{\prod_{i=1}^{\beta}(v_i - w)} \tag{2.15}$$

where v_i and w are respectively the rapidities of the holes and of the particle describing the intermediate state. Other form factors such that $\langle \lambda \mid \psi(x) \mid 0 \rangle$ confirm this description of the excitations. Although the description given here is reminiscent of a second quantized formalism, I am not aware that such a formalism exists.

3. THE SPIN CALOGERO-SUTHERLAND MODELS

A step towards the unification of the Calogero-Sutherland models and the integrable spin chains consists in introducing spin generalizations of the model studied in the last section.

The Hamiltonian we consider is

$$H = \sum_{j=1}^{N} \left(z_j \frac{\partial}{\partial z_j} \right)^2 - \sum_{i \neq j} \beta \left(\beta - P_{ij} \right) \frac{z_i z_j}{\left(z_i - z_j \right)^2} \tag{3.1}$$

The particles have an internal spin degree of freedom and the permutation P_{ij} exchanges the spins of the particles i and j.

The eigenstates have the following structure:

$$\psi \left(z_i, \sigma_i \right) = \Phi \left(z_i, \sigma_i \right) \prod_{i<j} \left(z_i - z_j \right)^{\beta} \tag{3.2}$$

where the wave function $\phi \left(z_i, \sigma_i \right)$ is completely antisymmetric under the simultaneous permutations of the spins and coordinates.

For our purpose, it is more convenient to look at the effective Hamiltonian acting on $\Phi \left(z_i, \sigma_i \right)$:

$$\tilde{H} = \sum_{j=1}^{N} \left(z_i \frac{\partial}{\partial z_j} \right)^2 + \beta \sum_{i \neq j} \frac{z_i + z_j}{z_i - z_j} \left(z_i \frac{\partial}{\partial z_i} - z_j \frac{\partial}{\partial z_j} \right) + \beta \sum_{i \neq j} \left(P_{ij} + 1 \right) \frac{z_i z_j}{\left(z_i - z_j \right)^2} \tag{3.3}$$

It turns out that this Hamiltonian can be diagonalized in a similar way as the one introduced in the last section. It has the same additive spectrum but the selection rules for the momenta differ from (2.10).

To do this, let us define the following Dunkl operators:

$$d_i = z_i \frac{\partial}{\partial z_i} - \beta i - \beta \sum_{j>i} \frac{z_i}{z_i - z_j} \left(K_{ij} - 1 \right) + \beta \sum_{j<i} \frac{z_j}{z_j - z_i} \left(K_{ij} - 1 \right) \tag{3.4}$$

where K_{ij} permutes the coordinates z_i and z_j :

$$\begin{array}{l} \left[K_{ij}, z_k \right] = 0 \quad \text{for} \quad k \neq i, j \\ K_{ij} z_j = z_i K_{ij} \end{array} \tag{3.5}$$

Some motivation for the form of these operators and the algebra which they obey will come later in the text (see also [17][18]), but first, let us see how they can be used to

solve the model. We shall show that the diagonalization of H can be reduced to the simultaneous diagonalization of the Dunkl operators d_i. Moreover in a convenient basis the d_i are represented by triangular matrices.

The d_i obey the defining relations of a degenerate affine-Hecke algebra:

$$\begin{aligned} &a) \quad [d_i, d_j] = 0 \\ &b) \quad [K_{ii+1}, d_k] = 0 \quad \text{if} \quad k \neq i, i+1 \\ &c) \quad K_{i,i+1} d_i - d_{i+1} K_{i,i+1} + \beta = 0 \end{aligned} \tag{3.6}$$

From these relations, one deduce that the quantity:

$$D(u) = \prod_{i=1}^{N} (u - d_i) = u^N + \sum_{k=1}^{N} c_k u^{N-k} \tag{3.7}$$

commutes with the permutations K_{ij}.

By computing explicitely $\sum d_i^2 = c_1^2 - 2c_2$, one sees that it is equal to the Hamiltonian $\tilde{H}$ except that the P_{ij} are replaced by $-K_{ij}$. Since $D(u)$ commutes with the permutations, we can restrict to the space of antisymmetric wave functions $\phi(z_i, \sigma_i)$. In this subspace $-K_{ij}$ and P_{ij} coincide and $\sum d_i^2$ is equal to $\tilde{H}$. Thus, instead of $\tilde{H}$, it is easier to simultanesouly diagonalize the d_i 's.

If we consider the explicit expression (3.4) of the d_i's, one sees that they preserve the space of homogeneous polynomials of a given degree in the variables z_i. It then follows from simple arguments that these operators are realized by triangular matrices in the monomial basis $z^\lambda = z_1^{\lambda_1} ... z_n^{\lambda_n}$ [3]. Their eigenvalues are equal (up to a permutation) to the components of the multiplet:

$$\delta_i = \lambda_i + \beta(N - i) \tag{3.8}$$

The eigenvectors of $\sum d_i^2$ corresponding to a given eigenvalue form a representation of the permutation algebra. The wave functions $\phi(z_i, \sigma_i)$ are obtained by antisymmetrizing these eigenvectors with an arbitrary spin function $X(\sigma_i)$. In particular, if we antisymmetrize the wave functions, we recover the Jack polynomials of the last section (at coupling $\beta + 1$) multiplied by the Vandermonde determinant $\prod_{i<j} (z_i - z_j)$. The mathematical properties of the Jack polynomials can probably be extanded to these wave-functions.

3.1. Dunkl operators and monodromy matrices

It is clear from the explicit construction of the wave functions of H that it's spectrum is highly degenerate and we shall now interpret these degeneracies from a symmetry principle. For this, we need to introduce the so-called monodromy matrix $T(u)$ solution of the equation:

$$R^{ab}T^aT^b = T^bT^aR^{ab} \tag{3.9}$$

where T^a in the operator valued matrix $T(u_a) \otimes 1$, T^b is $1 \otimes T(u_b)$ and R^{ab} is given by

$$R^{ab} = u_a - u_b + \beta P_{ab} \tag{3.10}$$

where P^{ab} permutes the two components of the tensor product $V^a \otimes V^b$.

Consistency of the above relations requires that R^{ab} is a solution of the Yang-Baxter equation in $V^a \otimes V^b \otimes V^c$:

$$R^{ab}R^{ac}R^{bc} = R^{bc}R^{ac}R^{ab} \tag{3.11}$$

Assume that $T(u)$ has a series expansion in $1/u$, the coefficients of this expansion define an algebra which is called the Yangian [19]. In this section, we exhibit a representation of the Yangian which commutes with $\tilde{H}$. For this, let us define the so-called "L operators "[20] defined as :

$$L_i = u - d_i - \beta P_{ai} \tag{3.12}$$

where P^{ai} permutes the components of the tensor product $V^a \otimes V^i$ and d_i is a coefficient which we identify with the Dunkl operator (3.4). This is consistent since the d_i's commute among themselves and with the permutations P_{ij}. A well known representation of the monodromy matrix $T(u)$ is then given by [20]:

$$T^a(u) = L_1(u)L_2(u)...L_N(u) \tag{3.13}$$

Now, let us act with this monodromy matrix on the space of wave functions $\phi(z_i, \sigma_i)$. The coordinates z_i are permuted by the K_{ij} and the spins σ_i are permuted by the P_{ij}. We shall see that the antisymmetric wave functions $\phi(z_i, \sigma_i)$ are preserved by the action of the monodromy matrix $T(u)$. To show this, we define a projection Π which substitutes $-P_{ij}$ for K_{ij} to the right of an expression.

The subspace of antisymmetric wave functions is preserved if the following condition is satisfied:

$$\Pi\left((K_{ij} + P_{ij})\,T(u)\right) = 0 \tag{3.14}$$

which is equivalent to the condition:

$$\Pi\left(\left(K_{ii+1}+P_{ii+1}\right)L_i(u)L_{i+1}(u)\right)=0 \tag{3.15}$$

when we expand this relation in powers of u, the coefficients of u and u^2 give relations which are trivially satisfied. Let us here consider the coefficient of 1. We omit the Π in front of the expression:

$$0=\left(K_{12}+P_{12}\right)\left(d_1d_2-\beta d_1P_{a2}-\beta d_2P_{a1}+\beta^2P_{a1}P_{a2}\right) \tag{3.16}$$

using (3.6) and the fact that $[K_{12},d_1d_2]=0$ we obtain:

$$\begin{aligned}-\beta P_{a1}P_{a2}P_{12}+\beta P_{12}P_{a1}P_{a2}\\ -d_2P_{12}P_{a1}-d_1P_{12}P_{a2}\\ +P_{a1}\left(d_1P_{12}-\beta\right)+P_{a2}\left(d_2P_{12}+\beta\right)=0\end{aligned} \tag{3.17}$$

which is easily shown. Notice that the unexpected term equal to β in (3.6)c) is necessary for the proof of this relation.

It follows from this analysis that the monodromy matrix $T(u)$ generates a representation of the Yangian algebra in the subspace of antisymmetric wave functions $\phi\left(z_i,\sigma_i\right)$. It is also easy to see that this representation commutes with the Hamiltonian $\tilde{H}$ and therefore, the Yangian is the symmetry algebra which explains the degeneracies of $\tilde{H}$.

3.2. Yang's representation of affine-Hecke algebras

In this section, we establish a relation between the Dunkl operators (3.4) and some operators defined by Yang in his study of the δ interacting gas (see also [21] for a discussion in the elliptic case).

To make this relation more transparent, we shall consider the case of the affine-Hecke algebra defined by :

$$\begin{aligned}&a)\quad [y_i,y_j]=0\\ &b)\quad [g_i,y_j]=0\quad\text{if}\quad j\neq i,i+1\\ &c)\quad g_iy_i-y_{i+1}g_i^{-1}=0\end{aligned} \tag{3.18}$$

and the g_i obey the relations of a Hecke algebra :

$$\begin{aligned}&[g_i,g_j]=0\quad\text{if}\quad |i-j|\geq 2\\ &g_ig_{i+1}g_i=g_{i+1}g_ig_{i+1}\\ &\left(g_i-q\right)\left(g_i+q^{-1}\right)=0\end{aligned} \tag{3.19}$$

This algebra replaces the permutations and the d_i's if one repeats the arguments of the last section replacing the rational solution (3.10) of the Yang-Baxter equation (3.11) with a trigonometric solution. This is considered in [3]. This algebra is also studied in the works of I.Cherednick [17]. Here we shall content ourselves to obtain a natural representation of the affine-Hecke algebra (3.18) which in some limit degenerates to the Dunkl operators (3.4).

For this, let us consider some realization of the Hecke-algebra (3.19) acting in the tensor product $V^{\otimes N}$ where the operator g_i acts in $V_i \otimes V_{i+1}$. To ease the forthcoming discussion, we rename $g_i : g_{i,i+1}$. Following Yang's argument [22], we shall exhibit a set of operators which commute together. Then we shall see that in some limit these operators can be identified with the y_j.

The starting point is the trigonometric form of the L operator [20] given by:

$$L_i^a(v) = \left(\frac{v}{v_i} g_{ia} - g_{ia}^{-1}\right) K_{ai} \tag{3.20}$$

where g_{ai} acts in $V_a \otimes V_i$ and K_{ai} permutes the two components of $V_a \otimes V_i$. Using this form we can construct a monodromy matrix solution of (3.9) given by :

$$T^a(u) = L_1(u) L_2(u) ... L_N(u) S^a \tag{3.21}$$

where S^a is a matrix acting in V^a and such that $[R^{ab}, S^a \otimes S^b] = 0$. It then follows from the Yang-Baxter equation (3.9) that the trace of this monodromy matrix $\bar{T}(v)$ defines a commuting set of transfer matrices:

$$\left[\bar{T}(v), \bar{T}(w)\right] = 0 \qquad \forall u,\ \forall v \tag{3.22}$$

To define the trace of the monodromy matrix, we consider T^a as a matrix acting in V^a and take its trace in the usual sense.

Hence, the operators $\tilde{y}_i = \bar{T}(v_i)$ form a commuting set . These operators play an important role in the study of the δ interacting gas [22]. They can be explicitely computed using the fact that:

$$L_i^a(v_i) = \left(q - q^{-1}\right) K_{ai} \tag{3.23}$$

If one then writes $\bar{T}(v_i)$ in the form $\bar{T}(v_i) = \text{tr } K_{ia}\theta$ where θ is an operator that contains only the permutations K_{ij} with $i, j \neq a$ and uses the following property of the trace:

$$\text{tr } K_{ai}\theta = \theta \tag{3.24}$$

one finally obtains :

$$\tilde{y}_i = X_{ii+1}X_{ii+2}...X_{iN}S^iX_{i1}...X_{ii-1} \tag{3.25}$$

where the operators X_{ij} are given by:

$$X_{ij} = \left(\frac{v_i}{v_j}g_{ij} - g_{ij}^{-1}\right)K_{ij} \tag{3.26}$$

The operators $\tilde{y}_i$ constructed in this way obey the relation (3.18)a) but not (3.18)b) c). It is then not difficult to see that these relations are satisfied in the limit $0 \ll v_1 \ll v_2... \ll v_N$. In this limit the operators y_j are given by [3][23]:

$$y_i = g_{ii+1}^{-1}K_{ii+1}...g_{iN}^{-1}K_{iN}S^iK_{1i}g_{1i}...K_{i-1,i}g_{i-1,i} \tag{3.27}$$

To make contact with the Dunkl operators d_i, we must use the following representation of the Hecke algebra acting in the space of polynomials in N variables z_i :

$$g_{ii+1} = qK_{ii+1} - (q-q^{-1})\frac{z_i}{z_i - z_{i+1}}(K_{ii+1} - 1) \tag{3.28}$$

and S^i is defined by:

$$(S^if)(z_1, ..., z_N) = f(z_1, ..., tz_i, ..., z_N) \tag{3.29}$$

If one sets $q = t^\beta$ and let $t \to 1$, the operators y_i tend to the d_i defined in (3.4).

4. THE HALDANE SHASTRY CHAIN

Among this class of models, the Haldane Shastry chain [7][8] is the closest to the known integrable spin chains such as the XXX chain [22]. It is also the less well understood. Here, we shall present these chains from the point of view of the Dunkl operators.

The Hamiltonian is the $\beta = \infty$ limit of (3.1), it is equal to :

$$H = \sum_{i<j}\frac{P_{ij}}{\sin^2(\theta_i - \theta_j)} \tag{4.1}$$

where the variables θ_j take the values $\frac{\pi j}{N}$, $0 \leq j < N$. The main difficulty comes from the fact that this Hamiltonian, unlike (3.1), cannot be obtained from the static limit of the Dunkl operators as $\sum d_i^2$ because in this limit this quantity is a number. In this section, we define a representation of the degenerate affine Hecke algebra (3.6) labelled by N complex numbers ω_k. When the ω_k are arbitrary, the representation is irreducible, when they are

equal to $\exp(i\frac{2\pi k}{N})$, it is reducible and one can find the Haldane Shastry Hamiltonian in its center. We shall then use a correspondence with the $\beta = 1$ representation (3.4) to diagonalise the Hamiltonian (4.1).

Consider the representation of the degenerate affine Hecke algebra (3.6) given by:

$$\tilde{d}_i = \sum_{j>i} \frac{z_i}{z_i - z_j} K_{ij} - \sum_{j<i} \frac{z_j}{z_j - z_i} K_{ij} \tag{4.2}$$

It acts on the Hilbert space defined by the permutations :

$$\mathcal{H}_N = \{|P_1 ... P_N > / P_i \in Z_N,\ P_i \neq P_j \quad \text{for all i, j}\} \tag{4.3}$$

in the following way :

$$\begin{aligned} K_{ij}|...P_i...P_j... > &= |...P_j...P_i... > \\ z_j|P_1...P_N > &= \omega_{P_j}|P_1...P_N > \end{aligned} \tag{4.4}$$

In the case where the ω_k's are arbitrary, there is no reason for this representation to be reducible. On the other hand, if $\omega_k = e^{i\frac{2\pi k}{N}}$, the operator C defined by:

$$C|P_j > = |(P_j + 1) > \tag{4.5}$$

obviously commutes with the d_j's and the K_{ij}. So, in this case the representation is reducible and one can verify that the Hamiltonian given by :

$$\tilde{H} = \sum_{i<j} \frac{z_i z_j}{(z_i - z_j)^2} K_{ij} \tag{4.6}$$

is also in the center. No systematic way to obtain the operators which commute with the K_{ij}'s and the d_i's is known. In what follows we restrict to the case where $\omega_k = \exp(i\frac{2\pi k}{N})$.

The Hamiltonian (4.6) can easilly be related to the Haldane Shastry Hamiltonian and we shall now indicate how to obtain its eigevalues and some of its eigenvectors. We shall construct a class of states on which the two representations of the d_i (3.4) (for $\beta = 1$) and (4.2) act in the same way. On these states, any conserved quantity of the spin chain has its analogue in the dynamical model and can thus be diagonalised .

For $M < N$ let us define the Hilbert space $\mathcal{H}_M$

$$\mathcal{H}_M = \{|P_1 ... P_M > / P_i \in Z_N,\ P_i \neq P_j \quad \text{for all i, j}\} \tag{4.7}$$

which is uniquely embedded in $\mathcal{H}_N$ if one requires that :

$$K_{ij}|P_1...P_M> = |P_1...P_M> \quad \text{for i, j} > \text{M} \tag{4.8}$$

In $\mathcal{H}_M$ we consider the following states :

$$|\alpha> = \Psi(\omega^{P_1},...,\omega^{P_M})|P_1,...,P_M> \tag{4.9}$$

where :

$$\Psi(z_i) = \Phi(z_i)\prod_{i<j}(z_i - z_j) \tag{4.10}$$

and $\Psi(z_i)$ is a polynomial of degree between 1 and $N-1$ in the variables z_i. On these states, $\tilde{d_i} = d_i - \frac{N+1}{2}$ for $1 \leq i \leq M$. It results from the fact that a polynomial $P(z)$ of degree between 1 and $N-1$ satisfies the relation :

$$(z\partial_z - \frac{N+1}{2})P(z)|_{z=\omega_k} = \sum_{l\neq k}\frac{\omega_k}{\omega_k - \omega_l}P(\omega_l) \tag{4.11}$$

The same relation enables to show that in this subspace $\tilde{H}$ acts as $\sum_{k=1}^{M}(d_k + 1/2 + N/2)(d_k + 1/2 - N/2)$ and to find its spectrum:

$$E = \sum_{k=1}^{M}\epsilon(f_k) \quad \text{with } \epsilon(\text{f}_\text{k}) = (\text{f}_\text{k} + 1/2)^2 - \text{N}^2/4 \tag{4.12}$$

$f_k + \frac{N+1}{2}$ are the degrees of the highest monomial of $\Psi(z_i)$. Therefore, the f_k satisfy the inequalities:

$$-\frac{N-1}{2} \leq f_1 \leq f_2... \leq \frac{N-1}{2} - 1 \tag{4.13}$$

References

[1] F.D.M. Haldane, To appear in "Proceedings of the 16th Taniguchi Symposium", 1993.

[2] C.F.Dunkl, Trans.Amer.Math.Soc, 311 (1989) 167.

[3] D.Bernard, M.Gaudin, F.D.M.Haldane and V.Pasquier, J.Phys.A 26 (1993) 5219.

[4] F.Lesage, V.Pasquier and D.Serban saclay preprint april 94.

[5] B. Sutherland, Phys.Rev.B 38 (1988), 6689.

[6] C.N.Yang Phys.rev.Letters, 19 (1967) 1312.

[7] F.D.M Haldane, Phys.Rev.Lett. 60 (1988) 635.

[8] B.S.Shastry, Phys.Rev.Lett. 60 (1988) 639.

[9] V.I.Inotzemtsev, J.Stat.Phys.59 (1990) 1143.

[10] I.V.Cherednick, preprint march 94.

[11] D.Bernard, V.Pasquier and D.Serban saclay preprint april 94.

[12] B.D. Simons, P.A. Lee, and B.L. Altshuler, Phys. Rev. Lett. V.70, No26, (1993) 4122.

[13] I.G. Macdonald, Séminaire Lotharingien, Publ. I.R.M.A. Strasbourg, 1988 .

[14] R. P. Stanley, Adv. in Math., 77, (1989) 76-115.

[15] F.D.M. Haldane, To appear in proceedings of the International Colloquium in Modern Field Theory, Tata institute, 1994.

[16] J. Minanan and A.P. Polychronackos CERN preprint april 94.

[17] I.V.Cherednick, Invent.Math.106 (1991) 411.

[18] A.P.Polychronakos,Phys.Rev.Letters, 69 (1992) 703.

[19] V.G.Drinfeld, Funct.Anal.Appl.20 (1988) 56.

[20] L.D.Faddeev Les Houches lectures, Elsevier Science Publishers (1984).

[21] V.M Buchstaber, G.Felder and A.P. Veselov preprint march 94.

[22] M.Gaudin, "la fonction d'onde de Bethe" Masson (1981).

[23] I.V.Cherednick, Commun.Math.Phys (1992) 150.